Automating Life Detection Using Lipid Detection in GCMS Data

P. Michael Furlong, Mary Beth Wilhelm, Denise Buckner, Morgan Anderson, Matt Chin, Tori Chinn, Tom McClure, Kanch Sridhar, Travis Boone, Jennifer Eigenbrode, Trey Smith, Tony Ricco, George Cooper, Linda Jahnke

9 February 2021










ExCALiBR


Extractor for Chemical Analysis of Lipid Biomarkers in Regolith


- 1. Building an instrument to extract and purify lipids from regolith to enable optimal GC-MS analysis.
- 2. Automatically classify lipids as being of *biotic* or *abiotic* origin.
- 3. Predict processing parameters for follow-on sampling and sample processing.

Gas Chromatography




Chromatogram





Mass Spectrometry







Image source: US Geological Survey via Wikipedia

Image source: US Government-NIST via H. Padleckas

Gas Chromatograph - Mass Spectrometer

- Combines both modalities to yield more information about samples.
- Amenable to operations on other planetary bodies.
- Good at detecting organic molecules that tell us about life (e.g. *lipids*)




Why Lipids?

Changes the question from:


Can organisms live here?

to:

Does this sample contain an organism?



Can We Classify Origin Process for Lipids?




- Not just looking for lipids in old places.
- Geological processes can transform signature of materials of *biotic* origin to look like materials of abiotic origin.

ABIOTIC


BIOTIC


Source: Assessing the diversity of lipids formed via Fischer-Tropsch-type reactions Mißbach et al., Organic Geochemistry 119 (2018)

Can We classify Origin Process for Lipids?




Preliminary Results: Yes






Leave-one-out validation results using K-Nearest Neighbour Classifier

More thrilling results!



Classification of 1931 biotic samples.

More thrilling results!



Classification of 1931 biotic samples.

Continuing Steps

- Improving Dataset quality
 - Existing dataset represents multiple decades of collection.
 - Abiotic samples are exceedingly rare.
- Considering alternative representations of GCMS data

